



# Recombinant Mouse TREM-1 (C-6His)

<b>Catalog #</b>	EPT118
<b>Expression Host</b>	Human Cells
<b>DESCRIPTION</b>	Recombinant Mouse Triggering Receptor Expressed On Myeloid Cells 1 is produced by our Mammalian expression system and the target gene encoding Ala21-Ser202 is expressed with a 6His tag at the C-terminus.
<b>Accession</b>	Q9JKE2
<b>Synonyms</b>	Triggering receptor expressed on myeloid cells 1; TREM-1; CD354; Trem1
<b>Mol Mass</b>	20.9 KDa
<b>AP Mol Mass</b>	38 KDa, reducing conditions
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
<b>FORMULATION</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.





## RECONSTITUTION

Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

It is not recommended to reconstitute to a concentration less than 100 $\mu$ g/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

## STORAGE

Lyophilized protein should be stored at  $< -20^{\circ}\text{C}$ , though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at  $4-7^{\circ}\text{C}$  for 2-7 days.

Aliquots of reconstituted samples are stable at  $< -20^{\circ}\text{C}$  for 3 months.

## BACKGROUND

Triggering Receptor Expressed on Myeloid Cells 1 (TREM-1) is a transmembrane protein with a single Ig-like domain. TREM-1 associates with the adapter protein, DAP12, to deliver an activating signal. TREM-1 is expressed on blood neutrophils and monocytes, and the expression is up-regulated by bacterial LPS.





TREM-1 is expressed at high levels on neutrophils of patients with microbial sepsis and in mice with a TREM-1/Fc fusion protein protected mice against LPS-induced shock. It stimulates neutrophil and monocyte-mediated inflammatory responses. Triggers release of pro-inflammatory chemokines and cytokines, as well as increased surface expression of cell activation markers. TREM-1 is amplifier of inflammatory responses that are triggered by bacterial and fungal infections and are a crucial mediator of septic shock.

## **SDS-PAGE**

