



# Recombinant Mouse IL-4 (C-6His)

|                        |   |
|------------------------|---|
| <b>Catalog #</b>       | EPT252  |
| <b>Expression Host</b> | Human Cells   |
| <b>DESCRIPTION</b>     | Recombinant Mouse Interleukin-4 is produced by our Mammalian expression system and the target gene encoding His21-Ser140 is expressed with a 6His tag at the C-terminus.        |
| <b>Accession</b>       | P07750  |
| <b>Synonyms</b>        | Interleukin-4; IL-4; IL4; B-cell IgG differentiation factor; B-cell growth factor 1; B-cell stimulatory factor 1; BSF-1; IGG1 induction factor; Lymphocyte stimulatory factor 1 |
| <b>Mol Mass</b>        | 14.6 KDa  |
| <b>AP Mol Mass</b>     | 15-19 KDa, reducing conditions  |
| <b>Purity</b>          | Greater than 95% as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>       | Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.  |
| <b>FORMULATION</b>     | Lyophilized from a 0.2 μm filtered solution of 20mM   |





PB, 150mM NaCl, pH 7.4.

## RECONSTITUTION

Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

It is not recommended to reconstitute to a concentration less than 100 $\mu$ g/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

## STORAGE

Lyophilized protein should be stored at  $< -20^{\circ}\text{C}$ , though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at  $4-7^{\circ}\text{C}$  for 2-7 days.

Aliquots of reconstituted samples are stable at  $< -20^{\circ}\text{C}$  for 3 months.

## BACKGROUND

Interleukin-4 (IL-4) is a pleiotropic cytokine that regulates diverse T and B cell responses including cell proliferation, survival and gene expression. IL-4 is produced by mast cells, T cells, and bone marrow stromal cells. IL-4 regulates the differentiation of naive





CD4+ T cells into helper Th2 cells, characterized by their cytokine-secretion profile that includes secretion of IL-4, IL-5, IL-6, IL-10, and IL-13, which favor a humoral immune response. Another dominant function of IL-4 is the regulation of immunoglobulin class switching to the IgG1 and IgE isotypes. Excessive IL-4 production by Th2 cells has been associated with elevated IgE production and allergic response.

## **SDS-PAGE**

